THE EFFECT OF LIVESTOCK GRAZING ON THE CHEMICAL PROPERTIES OF SOILS IN SUMMER LIVESTOCK CAMPS IN THE FANN MOUNTAINS (TAJIKISTAN)

Oimahmad Rahmonov [1], Michał Sobala [1], Katarzyna Pukowiec-Kurda [1], Buston Islamov [2], Zebiniso B. Islamova [3]

[1] Institute of Earth Sciences, University of Silesia in Katowice, Sosnowiec, Poland
[2] Department of Botany, Samarkand State University named after Sharof Rashidov, Samarkand, Uzbekistan
[3] Department of Medical Biology and Genetics, Samarkand State Medical University, Samarkand, Uzbekistan

Manuscript received: November 6, 2025
Revised version: January 21, 2026

Rahmonov O., Sobala M., Pukowiec-Kurda K., Islamov B., Islamova ZB. 2026. The effect of livestock grazing on the chemical properties of soils in summer livestock camps in the Fann Mountains (Tajikistan). *Quaestiones Geographicae* 45(3), Bogucki Wydawnictwo Naukowe, Poznań, pp. 69–88. 8 figs, 5 tables.

Abstract: High-mountain pastoral landscapes in Central Asia remain well preserved, providing an opportunity to assess the impact of pastoral activities on the natural environment. This study was conducted in the Fann Mountains (Tajikistan), where spring–summer grazing occurs. Areas where cattle rest during the day or night are known as qutans (summer cattle camps). This study aimed to analyse soil properties and their variability along a transect passing through summer cattle camps on geomorphologically diverse terrain. The studied soils are shallow and often lack their original humus horizon. On steep slopes, soil development is limited, whereas in low-slope areas, a thick organic layer accumulates due to animal excrement. Most soils were neutral to slightly alkaline in H_2O (pH 6.4–7.7), while pH in KCl ranged from 5.9 to 7.6. Organic carbon (TOC) and total nitrogen (TN) contents varied widely, ranging from 1.08% to 29.69% and 0.136% to 2.425%, respectively. Total phosphorus (TP) concentrations ranged from 201 mg·kg^{-1} to 13.520 mg·kg^{-1}, higher than the available phosphorus (AP: 4.36–4120 mg·kg^{-1}) for plants in all examined soil horizons. Elevated concentrations of TN, TOC and TP were observed in the central part of the qutans, where limited surface runoff and the accumulation of additional organic matter transported from higher elevations contribute to increased organic matter content in these areas.

Keywords: alpine pasture, sheep camping, grazing management, livestock grazing, soil chemistry

Corresponding author: Michał Sobala; michal.sobala@us.edu.pl

Introduction

Although mountains cover a relatively small proportion of the Earth's terrestrial surface, they are of great importance in terms of water resources, biodiversity, nutrient cycling and cultural values. Furthermore, they are characterised by a higher proportion of biodiversity than other areas (Körner et al. 2011). Mountain ecosystems are complex, dynamic, and exceptionally fragile, and are subject to global and local changes caused by human activity (Deléglise et al. 2019, Schmeller et al. 2022). Even in remote areas, human impacts can be significant, and these impacts, along with the resulting threats, remain largely understudied (Schmeller et al. 2018).

© 2026 Author(s)
This is an open access article distributed under the Creative Commons Attribution license

https://doi.org/10.14746/quageo-2026-0019
ISSN 2082-2103, eISSN 2081-6383

Mountain ecosystems are shaped by historical and contemporary land use practices (Körner et al. 1997, Lavorel et al. 2017), with the extent of influence depending mainly on the area's accessibility (Tasser, Tappeiner 2002, Jabs, Affek 2019, Rahmonov et al. 2024). Most of these changes are driven by pastoral activities, such as livestock grazing, which is the primary agricultural activity in most mountainous regions. Due to harsh natural conditions, other forms of human activity are often unsuitable. The diversity of natural and non-natural factors in different regions determines the development of various animal husbandry systems (Sobala 2014, Enchinov, Torushev 2022). As a result, the impact of livestock grazing on the environment varies across different mountain ranges.

In developed countries, pastoral activities in mountain regions have diminished due to socioeconomic factors and are now largely supported by subsidy policies (Mayer et al. 2022, Bucała-Hrabia 2024). By contrast, livestock grazing remains vital to the economies of developing countries, where it continues to shape forest ecosystems, soil health and the hydrologic cycle (Rahmonov et al. 2017a, Prokop et al. 2020).

Summer mountain pastures are intricate ecosystems where ecological processes and human activities are deeply interconnected. They provide essential fodder for domestic livestock during the summer and, through their traditional use, help sustain biodiversity, ecosystem services and open landscapes. Primarily used by local inhabitants, they also see some tourist activity. These pastures support local livelihoods as key areas for mowing and livestock grazing (Montenegro- Díaz et al. 2022).

In the high alpine grasslands of the Zarafshan mountains, pastoralism involves free grazing followed by a daily rest period of about 10 h. During this rest period, the animals gather in non-windy, flat slopes and low-lying areas, mainly near rivers and spring zones. These areas, known as summer qutans (summer camps), are used for grazing during this spring–summer period. They differ from the surrounding areas, creating a distinct spatial pattern marked by animal excrement. This leads to richer organic matter, which returns to the ecosystem in the form of manure and urine from the animals' consumption. By influencing grazing patterns and excreta deposition, the terrain's aspect affects soil nutrient concentration (Lambert, Roberts 1976).

The habitat conditions and soil characteristics of these areas remain underexplored and have received little attention in the literature. Some studies have partially addressed these factors in pastures in New Zealand (Sheath, Boom 1985, Haynes, Williams 1993), North America (Franzluebbers et al. 2000, Gill 2007), the Pyrenees (Aldezábal et al. 1999, Badía et al. 2008) and the Alps (Güsewell et al. 2005, Ferré et al. 2020). Nonetheless, the role of grazing animals in nutrient distribution within these ecosystems remains crucial. Grazing animals utilise only a small portion of the nutrients they consume, with 60–95% of the nutrients returning to the pasture as dung and urine, thereby influencing soil nutrient dynamics (Haynes, Williams 1999).

At the animals' resting sites, known as animal camps, dung accumulates to such an extent that vegetation is often absent. The soil in these areas becomes compacted and resistant to erosion. In the pasture, the recycling of nutrients from excreta back into the soil and plants affects the physical, chemical and biological properties of the soil (Iyyemperuma et al. 2007, Niu et al. 2009, Rahmonov et al. 2017a), as well as the vegetation composition. The impact of dung patches and urine on the chemical composition of the pasture and their influence on the soil under grazing conditions was analysed by Rodríguez et al. (1997, 2005). Their studies show that there was a significant effect of the dung patches and the micturitions on the pasture yield, both in the excretions deposited artificially and in those deposited directly by the cows in the grazing area. The highest values were produced with the micturitions, although in most of the cases the dung patches also caused significant increases. In turn, the effects of grazing itself were explored by Pineiro et al. (2010). Their studies show that identifying the pathway by which grazing may alter soil organic matter stocks in different regions is a key step for designing locally adapted management practices to increase soil organic matter storage.

The location of active cattle camps constantly shifts, while abandoned camps become overgrown with shrubs, leading to the formation of shrub thickets (Camacho et al. 2008, Rahmonov et al. 2017b, Losapio et al. 2024). Overgrazing combined with climate change – particularly extreme

weather events (Deléglise et al. 2019, Dong et al. 2020) – and unsustainable land management practices (Torresani et al. 2019) has led to changes in soil properties, including chemical and microbiological conditions (Maryskevych, Shpakivska 2008, 2011), vegetation structure and floral composition (Lyach, Milkina 2008, Jones et al. 2010). These changes result not only from the direct mechanical impact of cattle but also from natural fertilisation processes (Haynes, Williams 1999, Lyach, Milkina 2008). Nutrient and organic matter distribution and cycling in pasturelands are spatially heterogeneous, largely shaped by variations in the morphological conditions of the site (Haynes, Williams 1999, McIntire, Hik 2002). In mountain pastures, cattle typically seek out particular landscape features for resting and ruminating, such as areas beneath trees, shrubs, rocks, depressions and along watercourses (Rowarth et al. 1992, Haynes, Williams 1999).

While studies examining the influence of land use and natural processes on the functioning, physicochemical properties and structure of soil cover in the Central Asian Mountains are not uncommon (Abbasi et al. 2007, Wali et al. 2021, Zhao et al. 2013, Saeed et al. 2014, Rahmonov et al. 2017a), there is a notable lack of direct studies of soil chemistry in animal camping areas. As pastoral landscapes are predominantly preserved in Central Asia, this region offers an excellent opportunity to assess the impact of human activities on the environment. The study aims to assess the impact of grazing carried out in summer camps on the chemical properties of soils in mountain ecosystems of the Fann Mountains, Tajikistan. It was hypothesised that the chemical properties of the soils would vary both between different grazing areas (due to differences in grazing intensity) and within each grazing area (due to the migration of chemical components influenced by topographical variations). The main objective consists of specific goals: (1) to determine the differences in soil chemical properties between different camps due to varying grazing intensity and land use practices and (2) to identify the variability of chemical properties within the camps themselves, depending on topography.

Study area

The study area is located in the Fann Mountains (39°10'N, 68°15'E), which form the highest




Fig. 1. A – Study area Q1–Q7 summer livestock camp; B – an example of animal grazing on alpine meadows; C – Qutan in the Fann Mountains – a location in a terrain depression along a watercourse.

part of the Zarafshan Range in north-western Tajikistan (Figs 1 and 2). This range is part of the western part of Pamir-Alay (Rahmonov et al. 2013). The average altitude is approximately 4100 m a.s.l., with the highest peak being Chimtarga Peak at 5489 m a.s.l. The Fann Mountains feature typical high-altitude terrain, including U-shaped valleys, glacial cirques, steep slopes and numerous landslide cones (Rahmonov et al. 2013). The highest regions are covered by glaciers, while the valleys are home to glacial rivers with notable lakes such as Kuli Kalon, Alouddin, Dushakha and Iskandarkul, which possess significant geotourism potential (Pukowiec-Kurda et al. 2021). This mountain range is predominantly composed of rocks from the Palaeozoic (Carboniferous, Devonian, Silurian) and Mesozoic (mainly Cretaceous) eras, with younger Quaternary deposits filling the valley floors. The rocks include sandstones, quartzites, gneisses, limestones, marls, dolomites, as well as clays and loess. Detailed studies were conducted in the Urej River catchment (Fig. 1), where seven grazing areas (known as qutans) were selected for investigation: four located in the Kulikalon Basin and three near Artuch



Fig. 2. Photographs illustrating the study area – qutans in the Fann Mountains: A – Sheep grazing within the qutan Ovulgi Jalol (Q7-OJ) on a slope flattening in the Fann Mountains; B – A fragment of the qutan Murodali Sherali (Q2-MSh) in the headwater zone; C – Ovulgi Alisho sheep qutan (Q1-OA); D – Traces of summer livestock and the breeding of goats and cows during the summer period in Tushqoq (Q6-T); E – Dushacha camp near a water zone (Q4-D); F – Suchti Chuqurak unused cattle camp (Q5-SCh).

village. These areas are used for sheep grazing in the summer season (approximately 2 months) and vary in surface area, slope and distance from water sources (Table 1). Livestock grazing in the Fann Mountains occurs exclusively in summer, after which the herd gradually moves to the winter camps in the southern part of Tajikistan, near the Afghan border.

The diverse mosaic of habitats and topoclimatic conditions in the region supports a variety of plant formations that are vital to mountain ecosystems (Rahmonov et al. 2013) and provide valuable grazing areas for livestock. In Tajikistan, all mountain land is designated as grazing land by law (Law of the Republic of Tajikistan on Pasture 2013), encompassing vegetation types such as high mountain deserts, high mountain meadows and steppes; mid-mountain conifer and mesophytic forests; mid-mountain xerophytic light forests; mid-low-mountain semi-savannas; and wetlands. It is essential to highlight that Qutany and its surrounding areas are crucial habitats for *Biebersteinia multifida*, a medicinal plant utilised by local people for medicinal purposes. Used in folk medicine, this plant is believed to alleviate joint pain, reduce inflammation and soothe skin irritations. *Biebersteinia multifida* has medical significance and is traditionally used in Middle Eastern herbal medicine, with reported anti-inflammatory, analgesic and antioxidant effects, as well as anxiolytic (anti-anxiety) activity in animal studies (Zhang et al. 2020, Bustonovna 2025).

The valleys are dominated by grasslands and juniper forests, featuring *Juniperus seravschanica* and *Juniperus semiglobosa* (Rahmonov et al. 2017a). *J. semoglobosa* is found at higher elevations (2700–3200 m a.s.l.), forming juniper woodlands with varying density. *J. seravschanica* occurs at lower altitudes (1500–2400 m a.s.l.) (Rahmonov et al. 2024).

The climate and vegetation exhibit a distinct altitudinal zonation typical of this area. In the valley bottoms and lower slopes (1700–2000 m a.s.l.), July is the warmest month, with an average temperature of approximately 19°C, while January is the coldest month, with an average of about 4°C (Rahmonov et al. 2017a). At higher elevations (3000–3400 m a.s.l.), the average temperature in July drops to 10°C, and in January it falls to −10°C. Above the snow line (about 4000 m a.s.l.), the average annual temperatures are about −5°C, while at Chimtarga it can be as low as −15°C (Rahmonov et al. 2013). Mean yearly precipitation for the study area is 400–500 mm on peaks and slopes at altitudes of about 3000–3400 m a.s.l. (418 mm at Anzob Pass, 3373 m a.s.l.). In basins and deep valleys (altitudes in the range of

Table 1. Basic characteristics of summer cattle camps.

Summer cattle camp	Land use	Coordinates	Altitude	Slope	Length of transect	Vegetation
			[m a.s.l.]	[°]	[m]	
Q1-OA	Used	39°16'22"N 68°12'03"E	2935	9–17	35	*Galium aparine*, *Rumex Paulsenianus*, *Malva neglecta*, *Artemisia dracunculus*, *Convolvulus arvensis*, *R. Ecae*, *Berberis oblonga*
Q2-MSh	Used	39°16'14"N 68°11'59"E	2888	12–25	33	
Q3-B	Used	39°15'51"N 68°11'42"E	2913	16–19	40	
Q4-D	Used	39°15'02"N 68°11'50"E	2954	9–12	75	
Q5-SCh	Unused since 1989	39°20'09"N 68°07'43"E	2286	3–6	50	*Artemisia dracunculus*, *Convolvulus arvensis*, *Gentiana olivieri*, *Cichorium intybus*, *Salvia drobovii*, *Carex pachystilis*, *Arum Korolkovii*, *Plantago lanceolata*, *P. major*, *Iris songorica*, *Eremurusolgae*, *Ligularia thomsonii*, *Onosmadi chroanta*, *Centaurea squarrosa* and bushes as *Rosa Fedtschenkoana*, *R. huntica*, *R. Ecae*, *Berberis oblonga* and *Cerasus verrucosa*
Q6-T	Unused since 2015	39°19'38"N 68°08'02"E	2698	8–21	35	
Q7-OJ	Used	39°19'40"N 68°08'32"E	2851	4–19	85	*Galium aparine*, *Rumex Paulsenianus*, *Malva neglecta*, *Artemisia dracunculus*, *Convolvulus arvensis*, *R. Ecae*, *Berberis oblonga*

2200–2500 m a.s.l.), precipitation drops to 250–300 mm (271 mm in Iskanderkul, 2204 m a.s.l.). Most precipitation occurs in the spring season (60–70 mm in April or May), while winter is characterised by low rates of snowfall (Rahmonov et al. 2017a).

Materials and methods

Soil samples collection

The field study was conducted in the summer of 2019 and involved collecting soil samples for laboratory analysis.

Soil samples were collected from seven summer livestock camping areas across two parts of the Fann Mountains, within the Urej River catchment. These summer camps, known as ovul (or avul in some areas), serve as night-time resting sites for livestock. During the day, they are referred to as qutans (camps) and are typically set up in land hollows to keep animals grouped for protection against predators (Fig. 1C). The term 'summer livestock camps' used in this work refers to places where animals are grazed during the summer period. The analysed livestock camps can be categorised into two groups based on their primary purpose. The first group consists of summer camps used for breeding goats and cows during the summer months (Q5-SCh, Q6-T). The second group is dedicated to sheep grazing (Q1-OA, Q2-MSh, Q3-B, Q4-D, Q7-OJ) as part of transhumance – a seasonal movement of livestock between permanent summer and winter pastures.

Four of the summer livestock camp areas were located in the Kulikalon Basin and its immediate surroundings, while three were situated on the mountain slopes above the village of Artuch.

Six soil samples were collected from each campsite along the transect, from identified horizons, giving a total of 42 samples (Fig. 3).




Fig. 3. Schematic layout of transects and soil sampling sites within the summer livestock camp.

Samples were taken from the topsoil at different depths, depending on the degree of profile development or erosion. These included organic (O)/organic-mineral (OA) horizons, humus horizons (A) and cambic horizons (B).

In each case, the transect began at the upper boundary of the qutan, extended through its central part and ended at the lower boundary. Sampling was conducted from two depths in three sections of the qutan (Fig. 3):

- Upper part (IA and IB): Located on the slope between the mountain slope and the plateau. This area has sparse vegetation and shows signs of ongoing erosion;
- Central part (IIA and IIB): A flatter area where water flow slows down, allowing organic matter to accumulate;
- Lower part (IIIA and IIIB): Situated on the slope between the plateau and the lower area. This is the water runoff zone, though runoff is not always clearly visible.

Laboratory analysis

Forty-two soil samples were collected for laboratory analysis. The granulometric composition of the samples was determined using standard grain size analysis, which involved a sieve column with a set of sieves of varying mesh sizes. The following granulometric groups were distinguished: rock fragments (>10.0; 10.0–5.0; and 5.0–2.0 mm), fine earths (2.0–1.0; 1.0–0.5; 0.5–0.25; and 0.25–0.1 mm,) silt (0.1–0.05 mm) and clay (<0.05 mm). The mass of the sample remaining in each sieve was calculated as the percentage of grains of a given size in the total mass of the sample (Bednarek et al. 2004). In this paper, the results of the particle-size composition are presented as the sums of individual fraction groups.

The samples were air-dried, sieved (mesh sizes: 1 mm) and analysed as follows: pH was measured potentiometrically in both H_2O and 1N KCl using a glass electrode (soil:water:1/2.5); total organic carbon (TOC) was determined by the Tyurinn method; total nitrogen (TN) content was quantified using the Kjeldahl method; total phosphorus (TP) was measured by Bleck's method, modified by Gebhardt; available phosphorus (AP) was assessed using Egner–Riehm's method; and magnesium (AMg) was analysed according to PN-R-04023/23 (Bednarek et al. 2004).

Statistical analysis

To determine which chemical parameters differed between the samples, a difference test was performed. We analysed differences in chemical soil parameters between the upper and lower soil layers and differences in chemical parameters across the six types of soil samples (Fig. 3). Both analyses were conducted for the entire set of samples from all seven qutans. The dependent variables analysed were TOC, TN, TP, AP and AMg. Given the small sample size (n = 42), a non-parametric test was chosen.

To assess the differences in chemical soil parameters between the upper and lower soil layers, the Mann–Whitney test was applied. It is a non-parametric statistical test for determining differences between two groups on a single, ordinal variable with no specific distribution (Mann, Whitney 1947). To examine the differences in chemical parameters across the six types of soil samples, the Kruskal–Wallis test was used. The Kruskal–Wallis test is a non-parametric method for determining whether samples come from the same distribution (Kruskal, Wallis 1952). It is used to compare two or more independent samples, regardless of their sample sizes, and extends the Mann–Whitney test, which is used to compare only two groups.

It must be emphasised that each soil sample was treated as an independent observation; therefore, the results reflect differences between the individual samples rather than the entire qutans or qutan parts, and potential dependencies among samples from the same qutan were not accounted for. All analyses were performed using IBM SPSS Statistics version 31.0.0 (IBM Corp., Armonk, NY, USA), with a significance level of $\alpha = 0.05$. No correction for multiple comparisons was applied. To present the variation in soil chemistry across the transect segments and between the qutans, boxplots were used.

Results

Physical properties of soil

The analysed soil samples were taken from the topsoil, including the O, OA, A and enrichment (B) horizons. The general soil profile in the qutan areas is as follows: O-A-AC, OA-A-B and OB-B-C. The organic and organic-humus horizons are primarily composed of materials from animal excrement, mixed with mineral particles transported from higher areas due to erosion or surface runoff.

In terms of granulometric composition, fine earth fractions (73.9–91.9%; 2.0–0.1 mm), along with silt (3.4–14.1%, 0.1–0.05 mm) and clay (2–11.4%. <0.05), dominate all sites, except at the Q1-OA site, where rock fragments (>10–2 mm) are the dominant fraction, comprising 39.8% of the material (Fig. 4).

The pH values of the soil samples are generally close to neutral, with several samples even




Fig. 4. Granulometric composition of mineral horizons (Arabic numbers correspond to qutan numbers in Table 1).

being alkaline (Figs 5 and 6). This is primarily attributable to the geological substrate as well as the influence of grazing and animal excrement. The average pH values for the 42 samples were 7.14 in water and 6.83 in KCl, with ranges of 6.40–7.70 and 5.86–7.61, respectively.

Soil chemical properties

The basic descriptive statistics and normality test results are presented in Table 2. The Shapiro–Wilk test was conducted to assess the normality of the distribution for continuous variables. The results showed that the distributions for all chemical parameters deviated from normality ($p < 0.01$).

From the descriptive statistics, higher values are observed at the upper depth for all parameters (Table 2; Fig. 5). The highest values for the analysed chemical parameters were found in the middle part of the qutans (II). The only exceptions to this trend are TN and TOC, which reach their highest values at the upper depth in the lower part of qutan (IIIA). The coefficient of variation (CV) indicates very strong (AP, TP), strong (TOC in the lower depth and TN), medium (TOC in upper depth) and weak (AMg) variability in the parameter values.

Outliers were observed for all chemical parameters within qutan Q3-B (Fig. 6), particularly throughout the entire profile of the middle part (IIA, B) and the upper part of the profile in the lower section of the qutan (IIIA), where the results were significantly above the average values. Additionally, significantly higher values were found within qutan Q4-D for TN across the entire profile of the middle part (IIA, B), in the lower part of the profile in the upper section (IB) and the upper part of the profile in the lower section of the qutan (IIIA) (Fig. 6). For AMg, the elevated values were observed in the lower part of the profile in the upper section of the qutan (IB).

Phosphorus (P) content varied across the individual qutans. Both available P (4120 $mg \cdot kg^{-1}$) and total P (13520 $\cdot mg \cdot kg^{-1}$) were highest in the central parts of Q3-B and Q4-D, where the surface is flat, making these the most P-enriched values among all samples (Fig. 6).

TP shows less variability (Fig. 5) compared with AP. The highest concentrations of AP were also found in the Q4-D (1068 $mg \cdot kg^{-1}$) and Q1-OA (1262 $mg \cdot kg^{-1}$) sites, where TP also exhibited high values. However, high concentrations of TP do not affect AP concentrations in the Q2-MSh, Q5-Sch and Q7-OJ qutans. In these qutans, the highest AP values were generally found in the O and A horizons of the second sectors of the transects. In contrast, lower AP values were recorded in Q5-Sch (4.3676.30 $mg \cdot kg^{-1}$), Q7-OJ (33.14–119.46 $mg \cdot kg^{-1}$) and lower TP values in Q7-OJ (367–775 $mg \cdot kg^{-1}$) (Fig. 6).

Available magnesium concentrations varied across the individual qutans (Fig. 6). The average AMg values for the transect sections (I, IIB)

Table 2. Descriptive statistics grouped by chemical parameters and depth of soil sample collection, along with normality test results.

Parameter	Depth	Descriptive statistics								CV	W	p
		n	M	Mdn	Min	Max	IQR	SD	SK			
TN (%)	A	21	0.81	0.61	0.20	2.43	0.39	0.56	1.95	0.64	0.77	<0.01
	B	21	0.51	0.34	0.14	2.26	0.21	0.51	2.50	0.62	0.65	<0.01
TOC (%)	A	21	9.09	6.99	2.13	29.03	3.41	6.73	2.18	0.49	0.72	<0.01
	B	21	6.55	4.03	1.08	29.69	4.45	6.77	2.34	1.10	0.72	<0.01
AMg ($mg \cdot kg^{-1}$)	A	21	385.2	263.5	160.0	1752.5	71.0	435.4	2.91	0.27	0.46	<0.01
	B	21	346.8	214.0	132.0	1687.5	74.0	420.9	2.93	0.35	0.45	<0.01
AP ($mg \cdot P_2O_5 \cdot kg^{-1}$)	A	21	622.8	250.7	32.7	3684.2	744.2	875.9	2.53	2.97	0.68	<0.01
	B	21	518.1	76.3	4.4	4120.2	646.8	929.5	3.21	8.48	0.58	<0.01
TP ($mg \cdot kg^{-1}$)	A	21	2540.2	1885.0	224.8	12,980.0	1436.0	3140.7	2.70	0.76	0.61	<0.01
	B	21	1782.0	1112.0	201.2	13,520.0	1343.0	2858.6	3.79	1.21	0.52	<0.01

Annotation: n – subgroup size; AP – available phosphorus; CV – coefficient of variation; IQR – interquartile range; M – mean; Max – maximum value; Mdn – median; Min – minimal value; OA – organic-humus; P – *p*-value; SD – standard deviation; SK – Skewness; TN – total nitrogen; TOC – total organic carbon; TP – total phosphorus; W – Shapiro-Wilk's test statistics. A – stands for samples originating from the upper soil depth, representing the OA or humus (A) horizons; B – stands for samples collected from lower soil depth, located below the humus layers, mainly from the subsoil B (see Fig. 3).

of each qutan are as follows: Q3-B (990 mg·kg^{-1}), Q4-D (450 mg·kg^{-1}), Q5-Sch (254 mg·kg^{-1}), Q2 (241 mg·kg^{-1}), Q6-T (210 mg·kg^{-1}), Q1-OA (209 mg·kg^{-1}) and Q7-OJ (209 mg·kg^{-1}).

Similar trends were observed for organic carbon (TOC) and TN, as with phosphorus. The highest TOC concentrations were found in section IIB (Q3-B, 29.69%) and section IIIA (Q3-B, 29.03%). Similarly, the highest nitrogen content was recorded in the same sections, with values of 2.425% for Q3-B in section IIB and 2.258% for Q3-B in section IIIA. Additional differences in




Fig. 5. Variation of soil properties depending on the sampling location along the transect. Explanations: IA – upper part, upper depth; IB – upper part, lower depth; IIA – middle part, upper depth; IIB – middle part, lower depth; IIIA – lower part, upper depth; IIIB – lower part, lower depth (see Fig. 3). Explanation: The box represents the IQR, the horizontal line inside the box indicates the median, the cross (×) denotes the mean value and whiskers indicate the minimum and maximum values excluding outliers; outliers are shown as individual points.

TOC and TN concentrations are presented in Figure 5. The C/N ratio generally falls within a narrow range of 4–24 across most positions, except for Q6-T (C/N-53). These ranges are indicative of efficient organic matter metabolism by microorganisms.

Differences in chemical parameters

A non-parametric test (Mann–Whitney test) for independent samples revealed a statistically significant difference in the mean TN concentrations between the upper depth (A) and lower (B) soil layers. The mean TN in the upper layer (A)




Fig. 6. Variation of soil properties between individual qutans. Explanations: Q1-OA – Ovulgi Alisho, Q2-MSh – Murodali Sherali, Q3-B – Bibijon, Q4-D – Dushacha, Q5-SCh – Suchti Chuqurak, Q6-T – Tushqoq, Q7-OJ – Ovulgi Jalol (see Fig. 1). Explanation: The box represents the IQR, the horizontal line inside the box indicates the median, the cross (×) denotes the mean value and whiskers indicate the minimum and maximum values excluding outliers; outliers are shown as individual points.

was 0.812 (standard deviation [SD] = 0.563), while in the lower layer (B) it was 0.512 (SD = 0.513) (Table 3). Specifically, TN concentrations were significantly higher in the upper layer than in the lower layer.

A statistically significant difference for organic carbon (TOC) was found in the Mann–Whitney non-parametric test ($U = 100$; $p < 0.019$), with higher mean values in the upper depth (A) ($M = 9.089$; SD = 6.731) compared with the lower depth (B) ($M = 6.546$; SD = 6.769) (Table 3). For the remaining parameters, no significant differences were found between the upper and lower depths.

As far as differences in chemical parameters between qutan parts are concerned, a significant difference was found only for the TN parameter (Tables 4 and 5). Specifically, samples IB and IIIB exhibited significantly lower ranks compared with the other samples. This indicates that samples IB and IIIB share similar properties and are distinct from samples IA, IIA, IIIA and IIB,

Table 3. Results of difference test for upper and lower soil layer.

Parameter	Unit	Depth	Descriptive statistics		Mann–Whitney test			
			M	SD	Average rank	U	Z	*p*
Total nitrogen	[%]	A	0.81	0.56	27.24	100.0	−3.031	0.002
		B	0.51	0.51	15.76			
Total organic carbon	[%]	A	9.09	6.73	25.95	127.0	−2.352	0.019
		B	6.55	6.77	17.05			
Magnesium	[mg · kg^{-1}]	A	385.21	435.36	24.40	159.5	−1.532	0.125
		B	346.76	420.87	18.60			
Available phosphorus	[mg · P_2O_5 · kg^{-1}]	A	622.77	875.94	24.12	165.5	−1.385	0.166
		B	518.10	929.51	18.88			
Total phosphorus	[mg · kg^{-1}]	A	2540.28	3140.65	24.71	153.0	−1.698	0.090
		B	1781.98	2858.58	18.29			

Annotation: SD – standard deviation; A – stands for samples originating from the upper soil depth, representing the OA or humus (A) horizons; B – stands for samples collected from lower soil depth, located below the humus layers, mainly from the subsoil (B) (see Fig. 3).

Table 4. Descriptive statistics for six types of soil samples.

Parameter	Unit	Parameter	IA	IB	IIA	IIB	IIIA	IIIB	Total
Total nitrogen	[%]	M	0.49	0.39	1.01	0.80	0.93	0.35	0.66
		SD	0.19	0.42	0.69	0.73	0.61	0.16	0.55
		Min	0.20	0.14	0.49	0.27	0.43	0.21	0.14
		Max	0.74	1.32	2.43	2.26	2.18	0.66	2.43
Total organic carbon		M	5.85	5.47	10.79	9.66	10.62	4.51	7.82
		SD	1.95	4.95	7.63	9.71	8.40	4.06	6.79
		Min	2.13	1.22	4.08	2.64	4.80	1.08	1.08
		Max	7.66	15.30	25.84	29.69	29.03	13.24	29.70
AMg	[mg · kg^{-1}]	M	236.7	381.0	461.3	452.1	457.64	207.2	366.0
		SD	76.9	502.2	508.9	546.2	573.00	53.3	423.4
		Min	160.0	142.0	255.0	187.0	171.0	132.0	132.0
		Max	382.0	1517.5	1615.0	1687.5	1752.5	301.0	1752.5
Available phosphorus	[mg · P_2O_5 · kg^{-1}]	M	258.9	305.4	1050.7	1065.4	558.8	183.5	570.4
		SD	277.8	384.8	1261.4	1462.4	718.6	246.0	893.6
		Min	58.9	33.1	58.9	4.4	32.7	8.7	4.4
		Max	858.9	1068.2	3684.2	4120.2	2097.2	682.3	4120.2
Total phosphorus	[mg · kg^{-1}]	M	1224.1	1065.6	3631.7	3138.2	2765.0	1142.2	2161.1
		SD	856.4	1081.1	4228.0	4710.7	3286.4	810.5	2990.8
		Min	224.8	204.8	598.0	201.2	775.0	220.4	201.2
		Max	2605.0	3330.0	12,980.0	13,520.0	10,090.0	2455.0	13,520.0

SD – standard deviation.

Table 5. Results of Kruskal–Wallis test in six samples.

Sample parameter	Average rank				
	Total nitrogen	Total organic carbon	AMg	Available phosphorus	Total phosphorus
	[%]		[mg · kg^{-1}]	[mg · P_2O_5 · kg^{-1}]	[mg · kg^{-1}]
IA	21.00	21.07	18.07	20.43	19.14
IB	11.43	16.43	15.14	18.79	14.71
IIA	31.14	28.14	30.00	28.36	29.71
IIB	23.07	22.14	25.93	23.29	22.86
IIIA	29.57	28.64	25.14	23.57	25.29
IIIB	12.79	12.57	14.71	14.57	17.29
H Kruskal–Wallis	15.733	9.359	9.459	5.165	7.117
df	5	5	5	5	5
p	0.008	0.096	0.092	0.396	0.212

which are more similar to one another. No significant differences were observed for the other parameters.

Discussion

In this study, we demonstrated how soil chemical properties vary between different summer livestock camps and within each camp, depending on grazing intensity and topography. Our study examined the variability of soil properties along a transect within summer livestock camps (qutans). These camps differ in surface area, slope and proximity to water sources.

Summer livestock camps located above springs are characterised by steep surfaces and a mosaic of vegetation, whereas camps with even steeper slopes often have shallow soil profiles, with excrement deposited directly on the bedrock (Fig. 7A). This variation in terrain and camp characteristics likely leads to differences in soil chemical properties between camps. Erosion from cattle grazing and surface runoff leads to the accumulation of organic-mineral material in the central and lower sections of qutans, especially where steeper slopes cause greater runoff (Fig. 7B). The accumulation of organic matter in lower areas indicates that soil chemical properties also vary within individual camps depending on slope and runoff.

On the other hand, summer livestock camps with relatively gentle slopes (Q5-SCh) feature a thick organic horizon or its lack, primarily composed of animal-derived material. These areas are often almost devoid of plant cover, with only a single species growing in the shade of rocks, where the impact of animal fertilisation is minimal. Most of the summer livestock camps in the study area are located at altitudes between 2600 m a.s.l. and 3000 m a.s.l. (Rahmonov et al. 2017a). In contrast to the Kuli Kalon Basin, summer livestock camps near Artuch village are situated on brown mountain soils with thicker profiles, reaching depths of up to 1 m. These areas are characterised by abundant vegetation, which



Fig. 7. Contemporary usable qutan: located on a steep, eroded slope A – and accumulating organic-mineral materials as a result of erosion B.

is particularly vibrant in early spring and clearly marks the boundaries of the qutans, visible from distances of 10–15 km. In the Kuli Kalon Basin, the central areas of the summer camps appear mostly black, with narrow strips of vegetation marking the borders (Figs 2A, C, E).

The Urej River Basin, like other regions of the Fann Mountains, is characterised by significant elevation differences and high geomorphological diversity over relatively short distances. This variation contributes to a landscape with diverse terrain slopes and aspects, which in turn influence the range of vegetation, soil mosaic and the suitability of the area for summer livestock grazing, particularly goat and sheep breeding. Steep, high ridges and deep, narrow valleys foster soil erosion processes due to intensive water runoff, which often strips the soil cover down to the parent rock as a result of intensive grazing from historical times to the present (Rahmonov et al. 2017b, 2024). Consequently, most soil profiles in this area lack surface horizons, such as organic (O) and humus (A) layers. This situation has been observed mainly in unused qutans, such as Suchti Chuqurak (Q5-SCh), Tushqoq (Q6-T) and Ovulgi Jalol (Q7-OJ). In the case of Suchti Chuqurak (Q5-SCh), the area of qutans in 1987 was cultivated for barley by kolkhoz (Rahmonov et al. 2017a,b, 2024), which accelerated the soil erosion. The distinct physiographical features of the region play a key role in the formation of specific types of mountain soils (Kuteminskiy 1960, 1966); therefore, the occurrences of qutans in this region are also closely linked to terrain forms.

An important factor influencing soil properties is seasonal cattle grazing, which takes place from May to August and occurs both in stationary and transhumant systems, depending on climatic conditions. By the end of May, air temperatures in the Fan Mountains increase (to about 24°C) and with the end of the school year, the breeding and grazing season for domestic livestock begins. At the same time, sheep return from wintering grounds; in both cases, the grazing season ends in September. The terrain in the Urej River basin is highly diverse in geomorphological terms. The vast majority of the catchment area is characterised by slopes steeper than 40°, covering almost 98% of the basin. In contrast, the flattest surfaces (0–20°) occupy only 1.9 km^2 (0.7%) and are located in the Kulikalon Basin and in the vicinity of the village of Artuch (Rahmonov et al. 2024). Grazing on slopes steeper than 30° accelerates soil erosion and the degradation of soil horizons, thereby negatively affecting soil condition and stability. Additionally, livestock herds arrive at the summer livestock camp (qutans) both during the day and at night, where, as noted above, they remain for approximately 10 h. This results in the deposition of substantial amounts of animal excrement (dung and urine) into the soils of actively used qutans, including Ovulgi Alisho (Q1-OA), Murodali Sherali (Q2-MSh), Bibijon (Q3-B), Dushacha (Q4-D) and Ovulgi Jalol (Q7-OJ). The area of these camps typically ranges from 10 acres to 40 acres. In these areas, specific soil types form that are noticeably different from those found just beyond the borders of the qutan. This phenomenon has been observed in other mountain regions around the world, including cattle camps in New Zealand (Sheath, Boom 1985), the Pyrenees (Aldezábal et al. 1999) and the French Alps (Deléglise et al. 2019). Rahmonov et al. (2017a) also documented variations in soil chemistry relative to the distance from the qutan in the Fann Mountains.

The natural soils on which qutans are formed, according to the international classification (IUSS Working Group WRB 2022), include Calcaric Skeletic Fluvic Cambisol (Ochric) (Q3-B, Q4-D – lateral moraines), Calcaric Hyperskeletic Leptosol (Q1-OA, Q2-MSh – foot of the slope with active scree) and Cambisol (Q5-Sch, Q6-T, Q7-OJ – mountain slope). The position of the qutan within the terrain and the degree of slope inclination significantly influence the formation of the organic horizon and its distribution within the qutan.

In the studied qutans, notable differences were observed in the thickness of the organic-humus and humus horizons, as well as their enrichment (subsoil) in the individual studied cross-sections. The Q3-B area, and partly Q4-D, have relatively flat surfaces and are not subject to erosion, resulting in considerable thickness of the O and OA horizons. Organic matter from cattle appears in the form of a cover, creating a distinct boundary with the mineral horizons (A, AB), accumulating in layers.

In contrast, in section I of the transects, the horizons form as mixed organic-humus horizons (OA). The mineral material in these areas comes

from higher elevations, both due to grazing and surface runoff, which explains the dominance of the earth fraction (73.9–91.9%) in the analysed samples. Only in the Q1-OA site is the rock fragments fraction (>10–2 mm) dominant, comprising 39.8%. Similar results regarding soil horizon diversity and granulometric composition were observed at similar altitudes in the Pamir-Alay range (Tajikistan) by Rahmonov et al. (2017a) and Kabała et al. (2021).

In the Q5-Sch and Q7-OJ positions, horizon B reaches the surface directly due to the destruction of the humus horizon caused by overgrazing, the degree of slope and the susceptibility of the soil material to erosion. The fertility of qutan Q5-Sch led to its cultivation with wheat in 1988–1989, but the preceding ploughing further accelerated erosion, ultimately destroying the humus horizon.

Regarding granulometric composition, the analysed soil samples exhibit significant variability, particularly in the fraction >2.0 mm, which is most commonly associated with high altitude environments. In all samples, the fine fraction (2.0–0.1 mm) dominates within the humus horizon, partly due to material displacement caused by erosion from cattle near the qutans. This trend may also reflect the sampling focus, as only humus horizon samples were analysed. In this respect, the analysed soils differ from typical soils found in the upper mountain zone, which are characterised by high stoniness, a relatively coarse structure (low clay content), high carbonate levels, a neutral to alkaline reaction and low humus content (Kann 1965, Agakhanyantz, Lopatin 1978, Rahmonov et al. 2017a, Kabala et al. 2021). The analysed samples also exhibit a similar pH, largely due to the high carbonate content of the parent rock. Additionally, cattle excrement contributes to pH levels, further influencing soil chemistry.

The analysed soil horizons are rich in TN, organic carbon (TOC) and TP compared with neighbouring areas with less animal excrement. These elevated concentrations are closely related to cattle dung and urine. Higher values of TOC, TN and TP are found in almost all sections of IA (level OA), IIA and IIIB in the top layers, where animal excrement often forms a hard crust over the mineral layer. This crust, influenced by clay minerals, typically has low infiltration capacity, which results in lower concentrations of TOC, TN and TP in the deeper layers. It is important to note that plant communities are rarely present in qutan areas, limiting their contribution to the formation of soil organic matter.

The highest values of TOC, TN, AP, AMg and TP were recorded in Q3-B and Q4-D across almost all cross-sections and depths (Fig. 6). These findings confirm that grazing intensity and terrain morphology strongly influence nutrient accumulation in the soil. This is mainly due to the relatively flat surfaces of these areas, which are characterised by varied terrain morphology that systematically promotes the accumulation of organic matter from animal excrements, leading to the formation of a thicker organic horizon. In these areas, organic matter is not removed by surface runoff, which explains the thicker horizon and higher content of TOC, TN and TP. For example, in cross-section IIIB (Q4-D), the parent rock is located at a depth of 15 cm, with only the O or OA horizon above it. These results are also confirmed by similar TN content (Q3-B: IIa-2.425, 2.258%) and organic carbon (Q3-B: 29.69; 29.03%), despite varying depths – these levels show even distribution.

Similar findings for organic carbon (3.93–5.49%) and TN (0.50–0.84%) have been reported in studies of soil properties in livestock camps on summer pastures in the Pyrenees (Badía et al. 2008) and in sheep camps in Australia, where the C/N ratios were also comparable to those observed in the analysed area (Niu et al. 2009). Another possible explanation for these findings could be the random or point accumulation of specific factors. Similar patterns are observed in the distribution of magnesium and AP, where the highest values also occur in locations Q3-B and Q4-D. The reasons for this differentiation are similar to those of TOC, TN and TP.

Although the TP values are higher in these areas, they do not significantly impact the concentration of AP. Phosphorus, once released by mineral salts, quickly precipitates into less soluble forms, meaning only a portion of it remains available to plants. Additionally, low temperatures can further limit phosphorus availability. A drop in temperature below 13°C can reduce phosphorus availability by up to 70% (Li et al. 2020), which could help explain the low content of AP compared with TP.

Animal faeces contain more TP than AP because a substantial portion of phosphorus in

animal excreta occurs in organic forms, such as phytates, which are not directly available to plants (Barnett 1994, Pagliari, Laboski 2012). Organic phosphorus requires microbial mineralisation to become plant-available. This process is relatively slow and strongly dependent on environmental conditions, such as soil moisture and temperature, which may lead to a dominance of TP over AP in the studied soils. Areas of intensive grazing, where animals regularly deposit excrement, are often devoid of vegetation. Accumulated dung can form a compact crust on the soil surface, promoting surface runoff at the expense of water infiltration into deeper soil layers. Consequently, water retention is reduced, and microbial processes responsible for organic phosphorus mineralisation are slowed.

Soil temperature controls microbial activity and the rate of chemical reactions governing phosphorus cycling. Higher temperatures enhance microbial activity, accelerating the mineralisation of organic phosphorus (Mishra et al. 2015, Wu et al. 2019). In mountainous areas such as the Fann Mountains, extremely low or high temperatures may limit mineralisation and indirectly affect phosphorus availability by altering water retention and microbial conditions (Wu et al. 2019). Microorganisms can partially adjust their activity in response to temperature fluctuations, but this adaptation does not fully compensate for the constraints imposed by unfavourable thermal conditions (Bogati et al. 2024).

Soil pH plays a key role in regulating phosphorus availability (Amin 2023, Holland et al. 2024). Under acidic conditions, phosphorus is strongly bound by aluminium and iron oxides and hydroxides, reducing its availability to plants (Lemanowicz 2018, Huang et al. 2021). In alkaline soils, particularly those rich in calcium carbonate ($CaCO_3$), phosphorus can precipitate as sparingly soluble calcium phosphates, further limiting its availability (Xu et al. 2010, Wuenscher et al. 2015, Tandy et al. 2021). In practice, phosphorus availability is highest at near-neutral pH levels, whereas both low and high pH levels can restrict its plant accessibility.

Calcium carbonate primarily affects phosphorus dynamics by increasing pH and promoting the formation of insoluble calcium phosphates. High $CaCO_3$ concentrations in the studied soils (averaging 10–20% in the Fann Mountains; Rahmonov et al. 2017b) favour phosphorus precipitation, thereby limiting the pool of AP for plants. Variations in soil mineral composition and particle-size distribution further influence phosphorus retention and release (Lemanowicz, Krzyżaniak 2015). In slope soils, fine particles are often washed away, reducing the soil's water-holding capacity and consequently slowing the mineralisation of organic phosphorus. The measured soil pH in the qutans ranged from slightly acidic to slightly alkaline (6.4–7.7 in H_2O), generally near neutral, which is considered optimal for phosphorus availability. However, AP remained low in some qutan areas (Q2-MSh, Q5-SCh and Q7-OJ). The causes of low AP concentration in the case of Q2-MSh are related to the fresh inflow of animal excrement and its daily cleaning, which disrupts the natural processes of mineralisation of organic phosphorus substances. The soil in the Sukhti Chuqurak region (Q5-SCh) is largely devoid of organic matter due to past ploughing, and its loss has accelerated erosion processes that have affected the AP content. In the case of Ovulgi Jalol (Q7-OJ), the difference in AP compared with other analysed sites results from the degree of slope of the terrain and its intensive use, which translates into the lack of vegetation cover (apart from single species, Table 1).

In the vicinity of the studied qutans, across different plant communities, the results for AP and AMg ranged from 2.62 $mg \cdot kg^{-1}$ to 119 $mg \cdot kg^{-1}$ and 137 $mg \cdot kg^{-1}$ to 568 $mg \cdot kg^{-1}$, respectively. While the AP values are relatively low, they are comparable to the available magnesium concentrations observed in this study (Rahmonov et al. 2017a). It is important to note that only one sample in Rahmonov et al. (2017a) was taken from a qutan area, which featured an organic-humus horizon (OA), summer livestock camps, a very hard crust and a clearly layered surface with a 35° slope. The results from this location (39°15'41.37"N; 68°13'04.56"E) in terms of organic carbon (TOC) (11.4%), TN (1.072%), TP (9696 $mg \cdot kg^{-1}$) and available AP (1739 $mg \cdot kg^{-1}$) are similar to those found in Q3-B, Q4-D and Q6-T in some sections. This similarity further emphasises the role of animal herds in influencing soil chemistry. The lack of differentiation in some parameters can be attributed to the similarity in the composition of organic-humus, enriched soil horizons and the parent rock in these areas.

The role of animals in enriching the soil with phosphorus and nitrogen through their presence in various environments has been well documented (Rahmonov et al. 2014, Szymański et al. 2016). In open areas, animal excrement accelerates plant colonisation and soil-forming processes. However, in active qutans, these processes are inhibited, leading only to the formation of organic and humus horizons (Figs 2B and 8A). In contrast, in inactive camps (Q5-Sch), the soil's fertility facilitates the regeneration of the humus horizon that was previously degraded by grazing. In the Urej River Valley, mountain soils, where ecological systems are developing, are used for intensive summer grazing.




Fig. 8. Effects of using a summer livestock camp: qutan used in 2003 (A) and not used in 2012 – Tushqoq (Q6-T) (B), with bushes serving as potential resting areas for animals in Suchti Chuqurak (Q5-SCh) (C)

Grazing influences soil chemistry, but it also leads to the degradation of soil cover and its biological resources (Abbasi et al. 2007, Hagedorn et al. 2010). The mosaic distribution of organic matter is partly because, during the hottest part of the day (when temperatures reach 30°C), cattle seek cooler areas, such as the limits of the qutan, shrub canopies (Fig. 8) and the shade of rocks. This behaviour contributes to the high concentrations of TP and other soil properties in specific areas, a pattern also observed by other researchers (Metson, Saunders 1978, Pineiro et al. 2010, Ferré et al. 2020). As a result, large amounts of animal excrement accumulate in specific areas of the camp. The transfer of excrement to these areas represents a loss of nutrients from the surrounding land. Over time, the redistribution of nutrients within the pasture leads to an uneven distribution of soil fertility, which can disrupt the nutrient balance across the entire ecosystem (Haynes, Williams 1999, Güsewell et al. 2005).

Conclusions

Analyses were conducted to assess the formation of chemical properties in the soils of summer livestock camps, focusing on variations in chemical parameters along a transect that crosses the qutans and spans different soil depths. The results support the initial hypothesis that soil chemical properties vary both between qutans and within individual qutans, primarily due to topography and grazing patterns. The following conclusions can be drawn from the study:

The formation of the organic horizon in summer livestock camps depends on the topography and grazing intensity. Qutans with relatively flat surfaces and varied microtopography favour the accumulation of organic matter from animal excrement, resulting in the formation of a thick organic layer. Qutans with steep slopes, on the other hand, are more susceptible to surface runoff, which can wash away organic matter and, consequently, affect the overall soil chemistry.

Uneven deposition of animal excrements, resulting from the qutan topography and surface runoff, leads to the formation of so-called nutrient 'hot spots,' causing pronounced variability in soil chemical properties over relatively small areas within the qutan. Elevated concentrations of nitrogen, organic carbon and phosphorus are typically observed in the central parts of the qutan. In contrast, lower concentrations occur at their margins, which are more exposed to erosion caused by livestock trampling and surface runoff. Consequently, the chemical properties of soils within qutan areas differ significantly from those of soils in even the nearest neighbouring areas, forming so-called nutrient islands.

The chemical characteristics of soil in qutans vary depending on the location and whether the area is used for grazing. In the case of summer livestock camps that have been closed for economic reasons, continuous soil development processes occur. In areas permanently grazed, on the other hand, organic matter from animal excrement accumulates, and soil formation processes are partially inhibited due to the formation of a hard organic crust covering the soil.

Statistical analysis confirmed that TN and TOC are statistically significantly higher in the humus horizon (A) compared with the subsoil horizon (B).

Issues related to soil development within qutan areas and their immediate surroundings are of particular importance in the context of climate warming and carbon sequestration, especially since the qutans in Central Asia occupy extensive areas of mountainous regions.

Acknowledgements

We would like to thank the Reviewers for their careful reading of the manuscript and for their valuable comments and suggestions, which helped us to improve the quality of the paper

Authors' contributions

Conception and design: OR; Acquisition of data: OR, MS, KPK, BI and ZBI; Analysis and interpretation of data: OR, MS and ZBI; Drafting and/or revision of article: OR, MS and ZBI; Approval of the submitted version for publication: OR, MS, KPK, BI and ZBI.

References

Abbasi M.K., Zafar M., Khan S.R., 2007. Influence of different land-cover types on the changes of selected soil properties in the mountain region of Rawalakot Azad Jammu and Kashmir. *Nutrient Cycling in Agroecosystems* 78: 97–110. DOI 10.1007/s10705-006-9077-z.

Agakhanyantz O.E., Lopatin I.K., 1978. Main characteristics of the ecosystems of the Pamirs, USSR. *Arctic and Alpine Research* 10: 397–407. DOI 10.1080/00040851.1978.12003976.

Aldezábal A., Garin I., García-González R., 1999. Activity rhythms and the influence of some environmental variables on summer ungulate behaviour in Ordesa-Monte Perdido National Park. *Pirineos* 154: 145–157. DOI 10.3989/pirineos.1999.v153-154.110.

Amin A.E.A.Z., 2023. Effect of co-applying different nitrogen fertilizers with bone char on enhancing phosphorus release in calcium carbonate-rich soil: An incubation study. *Journal of Soil Science and Plant Nutrition* 23(2): 1565–1575. DOI 10.1007/s42729-023-01217-3.

Badía D., Fillat F., Martí C., Aguirre J., Gómez D., Sánchez J.R., 2008. Soil chemistry in long-term livestock camping areas in Pyrenean summer pastures. *Agrochimica* 52(3): 189–199.

Barnett G.M., 1994. Phosphorus forms in animal manure. *Bioresource Technology* 49(2): 139–147. DOI 10.1016/0960-8524(94)90077-9.

Bednarek R., Dziadowiec H., Pokojowska U., Prusinkiewicz Z., 2004. *Badania ekologiczno-gleboznawcze (Ecological and soil science research)*. PWN, Warsaw, Poland.

Bogati K., Sewerniak P., Walczak M., 2024. Impact of soil moisture on microbial diversity and their enzyme activity on agricultural soil. *bioRxiv*. DOI 10.1101/2024.10.07.617046.

Bucała-Hrabia A., 2024. Reflections on land use and land cover change under different socioeconomic regimes in the Polish Western Carpathians. *Regional Environmental Change* 24(1): 28. DOI 10.1007/s10113-024-02187-7.

Bustonovna I.Z., 2025. The yield of *Biebersteinia multifida* DC. Seeds in experiment and in nature. *Scientific Approach to the Modern Education System* 3(32): 108–111.

Camacho O., Dobremez L., Capillon A., 2008. Shrub encroachment in pastures in the Alps. Spatial organisation of farmers' activities and practices in Abondance valley (Haute-Savoie, France). *Journal of Alpine Research* 96(3): 89–100. DOI 10.4000/rga.575.

Deléglise C., Dodier H., Garde L., François H., Arpin I., Nettier B., 2019. A method for diagnosing summer mountain pastures' vulnerability to climate change, developed in the French Alps. *Mountain Research and Development* 39(2): D27–D41. DOI 10.1659/MRD-JOURNAL-D-18-00077.1.

Dong Z., Brahney J., Kang S., Elser J., Wei T., Jiao X., Shao Y., 2020. Aeolian dust transport, cycle and influences in high-elevation cryosphere of the Tibetan Plateau region: New evidence from alpine snow and ice. *Earth Science Reviews* 211: 103408. DOI 10.1016/j.earscirev.2020.103408.

Enchinov T., Torushev E., 2022. Summer camps of cattle breeders of the middle mountains of the Altai Republic. *Series Human Science* 4(2): 17–20. DOI 10.37882/2223-2982.2022.04-2.15

Ferré C., Caccianiga M., Zanzottera M., Comolli R., 2020. Soil–plant interactions in a pasture of the Italian Alps. *Journal of Plant Interactions* 15(1): 39–49. DOI 10.1080/17429145.2020.1738570.

Franzluebbers A.J., Stuedemann J.A., Schomberg H.H., 2000. Spatial distribution of soil carbon and nitrogen pools un-

der tall fescue. *Soil Science Society of America Journal* 64: 635–639. DOI 10.2136/sssaj2000.642635x.

Gill R.A., 2007. Influence of 90 years from grazing on plant and soil processes in the subalpine of the Wasatch Plateau, USA. *Rangeland Ecology and Managagement* 60: 88–98. DOI 10.2111/05-236R2.1.

Güsewell S., Jewell P.L., Edwards P.J., 2005. Effects of heterogeneous habitat use by cattle on nutrient availability and litter decomposition in soils of an alpine pasture. *Plant and Soil* 268: 135–149. DOI 10.1007/s11104-004-0304-6.

Hagedorn F., Mulder J., Jandl R., 2010. Mountain soils under a changing climate and land-use. *Biogeochemistry* 97: 1–5. DOI 10.1007/s10533-009-9386-9.

Haynes R.J., Williams P.H., 1993. Nutrient cycling and soil fertility in the grazed pasture ecosystem. *Advances in Agronomy* 49: 119–199. DOI 10.1016/S0065-2113(08)60794-4.

Haynes R., Williams P.H., 1999. Influence of stock camping behaviour on the soil microbiological and biochemical properties of grazed pastoral soils. *Biology and Fertility of Soils* 28(3): 253–258. DOI 10.1007/s003740050490.

Holland J.E., White R.P., Haefele S.M., Goulding K.W.T., McGrath S.P., Glendining M.J., 2024. Long-term effects of lime and phosphorus application on soil extractable (Olsen) phosphorus differ between two arable UK soils. *Soil Use and Management* 40(4): 1-9. DOI 10.1111/sum.13160.

Huang J., Zhou L.G., Liu S., Han T., Hayatu N.G., Li D., Zhang H., 2021. Vertical distribution of phosphorus fractions and the environmental critical phosphorus level in acidic red soil under long-term fertilizer and lime application in southern China. *Journal of Plant Nutrition and Soil Science* 184(5): 585–595. DOI 10.1002/jpln.202100098.

Iyyemperuma K., Israel D.W., Shi W., 2007. Soil microbial biomass, activity and potential nitrogen mineralization in a pasture: Impact of stock camping activity. *Soil Biology and Biochemistry* 39: 149–157. DOI 10.1016/j.soilbio.2006.07.002.

Jabs Z.J., Affek A., 2019. The agricultural accessibility of the Beskid Mountains. *Przegląd Geograficzny* 91(2): 97–111. DOI 10.7163/PrzG.2019.2.6.

Jones W.M., Fraser L.H., Curtis P.J., 2010. Plant community functional shifts in response to livestock grazing in intermountain depressional wetlands in British Columbia, Canada. *Biological Conservation* 143: 511–517. DOI 10.1016/j.biocon.2010.10.005.

Kabala C., Chachulski Ł, Gądek B., Korabiewski B., Mętrak M., Suska-Malawska M., 2021. Soil development and spatial differentiation in a glacial river valley under cold and extremely arid climate of East Pamir Mountains. *Science of the Total Environment* 758: 144308. DOI 10.1016/j.scitotenv.2020.144308.

Kann I.A., 1965. On the soils of the high altitude western Pamirs [in Russian]. *Pochvovedenie* 9: 16–25.

Körner W., Dupouey J.L., Dambrine E., Benoit M., 1997. Influence of past land use on the vegetation and soils of present-day forest in the Vosges Mountains, France. *Journal of Ecology* 85(3): 351–358. DOI 10.2307/2960507.

Körner C., Paulsen J., Spehn E.M., 2011. A definition of mountains and their bioclimatic belts for global comparisons of biodiversity data. *Alpine Botany* 121: 73–78. DOI 10.1007/s00035-011-0094-4.

Kruskal W.H., Wallis W.A., 1952. Use of ranks in one-criterion variance analysis. *Journal of the American Statistical Association* 47(260): 583–621. DOI 10.1080/01621459.1952.10483441

Kuteminskiy V.J., 1960. *About soils of Pamir*. Tajik SSR Academy of Science Press, Dushanbe, Tajikistan.

Kuteminskiy V.J., Leontyeva R.S., 1966. *The soils of Tajikistan*. Irfon, Dushanbe, Tajikistan.

Lambert M.G., Roberts E., 1976. Aspect differences in an unimproved hill country pasture. 1. Climatic differences. *New Zealand Journal of Agricultural Research* 19(4): 459–467. DOI 10.1080/00288233.1976.10420975.

Lavorel S., Grigulis K., Leitinger G., Kohler M., Schirpke U., Tappeiner U., 2017. Historical trajectories in land use pattern and grassland ecosystem services in two European alpine landscapes. *Regional Environmental Change* 17: 2251–2264. DOI 10.1007/s10113-017-1207-4.

Law of the Republic of Tajikistan, 2013. On pasture. Law No. 951, Article 194, Dushanbe, Mar 19, 2013. Online: majmili.tj (accessed 17 January 2025).

Lemanowicz J., 2018. Dynamics of phosphorus content and the activity of phosphatase in forest soil in the sustained nitrogen compounds emissions zone. *Environmental Science and Pollution Research* 25(33): 33773–33782. DOI 10.1007/s11356-018-3348-5.

Lemanowicz J., Krzyżaniak M., 2015. Vertical distribution of phosphorus concentrations, phosphatase activity and further soil chemical properties in salt-affected Mollic Gleysols in Poland. *Environmental Earth Sciences* 74(3), 2719-2728. DOI 10.1007/s12665-015-4294-x.

Li F.Y., Yuan C.Y., Yuan Z.Q., You Y.J., Hu X.F., Wang S., Li G.Y., 2020. Bioavailable phosphorus distribution in alpine meadow soil is affected by topography in the Tian Shan Mountains. *Journal of Mountain Science* 17(2): 410–422. DOI 10.1007/s11629-019-5705-3.

Losapio G., De Moraes C.M., Nickels V., Tscheulin T., Zouros N., Mescher M.C., 2024. The effects of shrub encroachment on arthropod communities depend on grazing history. *Global Ecology and Conservation* 50: e02819. DOI 10.1016/j.gecco.2024.e02819.

Lyach I.V., Milkina L.I., 2008. Nacional'nyi pryrodnyi park "Skolivski Beskydy". Nelisowaroslynnist' (National Nature Park "Skolivski Beskydy". Vegetation). Spolom, Lviv, Ukraine.

Mann H.B., Whitney D.R., 1947. On a test of whether one of two random variables is stochastically larger than the other. *The Annals of Mathematical Statistics* 18: 50–60. DOI 10.1214/aoms/1177730491.

Maryskevych O., Shpakivska I., 2008. Fermentativna aktyvnist' gruntiv v ekosystemach Schidnich Beskydiv Ukrains'kich Karpat (Enzymatic activity of soils in the ecosystems of the Eastern Beskids of the Ukrainian Carpathians). In: *Geneza, geografia ta ekologia gruntiv. Zbirnik naukovych prac na poshanu profesora LNU im. Ivana Franka Myrona Kita*. Vydavnychij centr LNU im, Ivana Franka, Lviv, Ukraine: 342–347.

Maryskevych O., Shpakivska I., 2011. Wpływ użytkowania pasterskiego na właściwości gleb w Beskidach Skolskich (ukraińska część Karpat Wschodnich) (Effect of pastoral use on soil properties in the Skole Beskids, Ukrainian part of the Eastern Carpathians). *Roczniki Bieszczadzkie* 19: 349–358.

Mayer A., Egger C., Loyau A., Plutzar C., Schmeller D.S., Gaube V., 2022. Mountain pastures increase the resilience of livestock farming to extreme events in the Ariège department, France. *Agronomy for Sustainaible Development* 42: 49. DOI 10.1007/s13593-022-00779-3.

McIntire E.J., Hik D.S., 2002. Grazing history versus current grazing: leaf demography and compensatory growth of three alpine plants in response to a native herbivore (Ochotona collaris). *Journal of Ecology* 90(2), 348-359. DOI 10.1046/j.1365-2745.2001.00672.x.

Metson A.J., Saunders W.M.H., 1978. Seasonal variations in chemical composition of pasture. *New Zealand Journal of Agricultural Research* 21(2): 341–353. DOI 10.1080/00288233.1978.10427419.

Mishra A., Pattnaik T., Das D.N., Das M., 2015. Vertical distribution of available plant nutrients in soils of mid central valley at Odisha zone, India. *American Journal of Experimental Agriculture* 7(4): 214–221. DOI 10.9734/ajea/2015/15611.

Montenegro-Díaz P., Alvear R.C., Wilcox B.P., Carrillo-Rojas G., 2022. Effects of heavy grazing on the microclimate of a humid grassland mountain ecosystem: Insights from a biomass removal experiment. *Science of the Total Environment* 832: 155010. DOI 10.1016/j.scitotenv.2022.155010.

Niu Y., Li G.D., Li L., Chan K.Y., Oates A., 2009. Sheep camping influences soil properties and pasture production in an acidic soil of New South Wales, Australia. *Canadian Journal of Soil Science* 89: 235–244. DOI 10.4141/CJSS0800.

Pagliari P.H., Laboski C.A., 2012. Investigation of the inorganic and organic phosphorus forms in animal manure. *Journal of Environmental Quality* 41(3): 901–910. DOI 10.2134/jeq2011.0451.

Pineiro G., Paruelo J.M., Oesterheld M., Jobbágy E.G., 2010. An assessment of grazing effects on soil carbon stocks in grasslands. *Rangeland Ecology and Management* 63(1): 109–119. DOI 10.2111/08-255.1.

Prokop P., Wiejaczka Ł, Sarkar S., Bryndal T., Bucała-Hrabia A., Kroczak R., Soja R., Płaczkowska E., 2020. Morphological and sedimentological responses of small stream channels to extreme rainfall and land use in the Darjeeling Himalayas. *Catena* 188: 104444. DOI 10.1016/j.catena.2019.104444.

Pukowiec-Kurda K., Rahmonov O., Sobala M., Myga-Piątek U., 2021. The assessment of hydrogeosites in the Fann Mountains, Tajikistan as a basis for sustainable tourism. *Resources* 10(12): 126. DOI 10.3390/resources10120126.

Rahmonov O., Holbegov M., Szczypek T., Snytko V.A., Kłys G., Rahmonova M., 2014. The consequences of vegetation degradation under the influence of anthropogenic activity in the territory of the Zarafshan Range (Western Tajikistan). *Geography and Natural Resources* 3: 193–197. DOI 10.1134/S1875372814020127.

Rahmonov O., Majgier L., Andrejczuk W., Banaszek J., Karkosz D., Parusel T., Szymczyk A., 2013. Landscape diversity and biodiversity of Fann Mountains (Tajikistan). *Ekológia* 32: 388–395. DOI 10.2478/eko-2023-0037.

Rahmonov O., Rahmonov M., Opała-Owczarek M., Owczarek P., Niedźwiedź T., Myga-Piątek U., 2017a. Ecological and cultural importance of juniper ecosystem in the area of Zerafshan Valley (Tajikistan) on the background of environmental condition and anthropogenic hazards. *Geographia Polonica* 90: 441–461. DOI 10.7163/GPol.0110.

Rahmonov O., Szczypek T., Niedźwiedź T., Myga-Piątek U., Rahmonov M., Snytko V.A., 2017b. The human impact on the transformation of juniper forest landscape in the western part of the Pamir-Alay range (Tajikistan). *Environmental Earth Sciences* 76: 1–17. DOI 10.1007/s12665-017-6643-4.

Rahmonov O., Szypuła B., Sobala M., Islamova Z.B., 2024. Environmental and land-use changes as a consequence of land reform in the Urej River catchment (Western Tajikistan). *Resources* 13(4): 59. DOI 10.3390/resources13040059.

Rodríguez I., Crespo G., 1997. Effect of cattle dung and urine on yield and N, P and K content of *Cynodon nlemfuensis* pasture. *Cuban Journal of Agricultural Science* 31: 183-188.

Rodríguez I., Crespo G., Torres V., Fraga S., 2005. Effect of the dung patches and the urine on the chemical composition of the pasture and their effect on the soil under grazing conditions or not. *Cuban Journal of Agricultural Science* 39(3): 341–350.

Rowarth J.S., Tillman R.W., Gillingham A.G., Gregg P.E.H., 1992. Phosphorus balances in grazed, hill-country pastures: The effect of slope and fertiliser input. *New Zealand Journal of Agricultural Research* 35(3): 337–342. DOI 10.1080/00288233.1992.10427511.

Saeed S., Younus M., Barozai K., Ahmad A., Shah S.H., 2014. Impact of altitude on soil physical and chemical properties in Sra Ghurgai (Takatu mountain range) Quetta, Balochistan. *International Journal of Science and Engineering Research* 5(3): 730-735.

Schmeller D.S., Loyau A., Bao K., Brack W., Chatzinotas A., De Vleeschouwer F., Friesesn J., Gandois L., Hansson S.V., Haver M., Le Roux G., Shen J., Teisserenc R., Vredenburd V.T., 2018. People, pollution and pathogens – global change impacts in mountain freshwater ecosystems. *Science of the Total Environment* 62(2–623): 756–763. DOI 10.1016/j.scitotenv.2017.12.006.

Schmeller D.S., Urbach D., Bates K., Catalan J., Cogălniceanu D., Fisher M.C., Friesen J., Füreder L., Gaube V., Haver M., Jacobsen D., 2022. Scientists' warning of threats to mountains. *Science of the Total Environment* 853: 158611. DOI 10.1016/j.scitotenv.2022.158611.

Sheath G.W., Boom R.C., 1985. Effects of November–April grazing pressure on hill country pastures 3. Interrelationship with soil and pasture variation. *New Zealand Journal of Experimental Agriculture* 13(4): 341–349. DOI 10.1080/03015521.1985.10426102.

Sobala M., 2014. Krajobrazy pasterskie w Polsce i Europie – wybrane typy, przykłady i formy ich ochrony (Pastoral landscapes in Poland and Europe – selected types, examples and forms of protection). *Prace Komisji Krajobrazu Kulturowego* 25: 81–98.

Szymański W., Wojtuń B., Stolarczyk M., Siwek J., Waścińska J., 2016. Organic carbon and nutrients (N, P) in surface soil horizons in a non-glaciated catchment, SW Spitsbergen. *Polish Polar Research* 37: 49–66. DOI 10.1515/popore-2016-0006.

Tandy S., Hawkins J.M.B., Dunham S.J., Hernández-Allica J., Granger S.J., Yuan H., Blackwell M.S.A., 2021. Investigation of the soil properties that affect Olsen P critical values in different soil types and impact on P fertiliser recommendations. *European Journal of Soil Science* 72(4): 1802–1816. DOI 10.1111/ejss.13082.

Tasser E., Tappeiner U., 2002. Impact of land use changes on mountain vegetation. *Applied Vegetation Science* 5: 173–184. DOI 10.1111/j.1654-109X.2002.tb00547.x.

Torresani L., Wu J., Masin R., Penasa M., Tarolli P., 2019. Estimating soil degradation in montane grasslands of North-eastern Italian Alps (Italy). *Heliyon* 5(6): e01825. DOI 10.1016/j.heliyon.2019.e01825.

Wali E., Tasumi M., Shinohara Y., 2021. Classification and digital mapping of soils in a semiarid region of Afghanistan. *Eurasian Soil Science* 54: 38–48. DOI 10.1134/S1064229321010142.

Working I.U.S.S., Group W.R.B., 2022. World Reference Base for Soil Resources. International soil classification system for naming soils and creating legends for soil maps, 4th Edn. International Union of Soil Sciences (IUSS), Vienna, Austria.

Wuenscher R., Unterfrauner H., Peticzka R., Zehetner F., 2015. A comparison of 14 soil phosphorus extraction

methods applied to 50 agricultural soils from Central Europe. *Plant, Soil and Environment/Czech Academy of Agricultural Sciences* 61(2): 86–96. DOI 10.17221/932/2014-pse.

Wu Y., Wang Q., Wang H., Wang W., Han S., 2019. Shelterbelt poplar forests induced soil changes in deep soil profiles and climates contributed their inter-site variations in dryland regions, northeastern China. *Frontiers in Plant Science* 10: 220. DOI 10.3389/fpls.2019.00220.

Xu W., Chen X., Luo G., Zhang Q., Qing L., 2010. Soil properties at the tree limits of the coniferous forest in response to varying environmental conditions in the Tianshan Mountains, northwest China. *Environmental Earth Sciences* 63(4): 741–750. DOI 10.1007/s12665-010-0745-6.

Zhang B., Jin X., Yin H., Zhang D., Zhou H., Zhang X., Tran L.S.P., 2020. Natural products, traditional uses and pharmacological activities of the genus *Biebersteinia* (Biebersteiniaceae). *Plants* 9(5): 595. DOI 10.3390/plants9050595.

Zhao Z., Zeng H., Wu J., Zhang L., 2013. Organochlorine pesticide (OCP) residues in mountain soils from Tajikistan. *Environmental Science Processes and Impacts* 15(3): 608–616. DOI 10.1039/c2em30849e.